### California Global Warming Solutions Act of 2006

# **Greenhouse Gas Inventory and 1990 Emissions Level**

### Implementation of AB 32 Requirements

California Air Resources Board August 13, 2007 - Cal/EPA Headquarters

### Workshop Agenda

- Greenhouse Gas Inventory
- Database and Documentation
- 1990 Emissions Level
- 2020 Projections Ongoing Work
- Schedule



### Background

## AB 32 Emission Inventory Activities

- Transfer from CEC to ARB on January 1, 2007
- Update statewide GHG Inventory
- Determine statewide 1990 emissions level
- Establish statewide 2020 emissions limit equal to 1990 emissions level
- Present to Board by January 1, 2008

# Description of Statewide Inventory

- Emissions of six "Kyoto gases" in tonnes of CO2 equivalents (CO2e)
- Generally consistent with IPCC and U.S.EPA guidance
- Uses IPCC Second Assessment Report GWPs
- "Top-down" vs. "bottom-up" approaches


### Inventory Updates

- Improved data input
  - Updated emission factors
  - Improved activity data
  - Corrected calculation errors
- Methodological improvements for some categories
- Data sources:
  - International organizations
  - Federal and state agencies
  - Stakeholders


### Scope of Inventory Review

- All sectors and ~270 sub-categories
- More than 10,000 calculations
- Entire inventory for 1990-2004


# 1990 Gross Emissions by Fuel (MMTCO2 Eq)



# 1990 Gross Emissions by Gas (MMTCO2 Eq)



## Updated 1990 Gross Emissions



# Results of Review (MMTCO2 Eq)

Sector	1990 CEC	1990 ARB	% difference
Agriculture	27	22	-19%
Cement	5	8	60%
Electricity Generation – In-State	45	47	4%
Electricity Generation – Imported	44	53	20%
Landfills	8	7	-13%
Refineries	28	30	7%
Land Use, Land Use Change and Forestry (net, biodegradable CO2)	-17	-6	-65%
Transportation	178	171	-4%
Total (all sectors)	413	436	



### Sector Technical Discussion

### Focus of Inventory Review

- Agriculture
- Cement
- Electricity Generation
- Refineries
- Landfills
- Land Use, Land Use Change & Forestry
- Transportation

~80% of Total 1990 Emissions

### Agriculture


- Includes:
  - Manure Management
  - Enteric Fermentation
  - Agricultural Residue Burning
  - Managed Soils

#### Manure Management - Updates

- Includes emissions from management of livestock waste
- Data Sources: California-specific data compiled by U.S.EPA
- Updates:
  - Used 2006 IPCC methodology
  - Used data from U.S.EPA 1990-2005 inventory

### Manure Management - Estimates

CEC 1990 estimate:4.10 MMTCO2 Eq



New 1990 estimate:5.00 MMTCO2 Eq

#### **Enteric Fermentation - Updates**

- Includes exhaled methane emissions that are produced in animals' digestive systems
- Data Sources: California-specific data compiled by U.S.EPA

- Update:
  - Used most recent U.S.EPA model outputs

### Enteric Fermentation - Estimates

CEC 1990 estimate:7.53 MMTCO2 Eq

New 1990 estimate:6.67 MMTCO2 Eq



## Agricultural Residue Burning - Updates

- Includes emissions (CH4 and N2O) from burning Ag biomass waste
- Data Sources: California-specific data compiled by CEC

- Updates:
  - Accounted for rice straw phase-down
  - Revised calculations

### Agricultural Residue Burning – Estimates

CEC 1990 estimate:0.14 MMTCO2 Eq

New 1990 estimate: 0.33 MMTCO2 Eq



### Managed Soils - Updates

- Includes Ag practices that add nitrogen to soil and increase N2O emissions
- Data Sources: California-specific data compiled by CEC

- Updates:
  - Revised calculations
  - Improved emission factors

### Managed Soils - Estimates

CEC 1990 estimate: 14.68 MMTCO2 Eq

New 1990 estimate: 9.52 MMTCO2 Eq



#### Cement

Includes emissions from clinker production and fuel use

- Data Sources:
  - USGS statewide clinker production data
  - California Energy Balance
  - IPCC emission factors


#### Cement - Updates

- Updated clinker production data
- Used IPCC 2006 Revised Guidelines methodology for process CO2 emissions
- Included 5 year average for coke and coal

#### Cement - Estimates

CEC 1990 estimate:4.80 MMTCO2 Eq

New 1990 estimate:8.23 MMTCO2 Eq



### **Electricity Generation**

Includes emissions from generating in-state and imported electricity

- Data sources:
  - Energy Information Administration
  - IPCC emission factors for CH4 and N2O

## In-State Electricity Generation - Updates

- Included additional fuels (i.e., biomass, geothermal)
- Used IPCC oxidation factor of 1
- Updated SF6 emissions
- Included CHP useful thermal output in industrial and commercial sector emissions

## Imported Electricity Generation - Updates

- Included 6 additional specified imports and updated 2 specified imports
- Used gross imported power
- Accounted for line loss
- Included IPCC emission factors for CH4 and N2O
- Used IPCC oxidation factor of 1
- Updated SF6 emissions

#### Electricity Generation - Estimates

- In-State CEC 1990 estimate: 45.17 MMTCO2 Eq
- In-State New 1990 estimate: 46.95 MMTCO2 Eq
- Imported CEC 1990 estimate: 44.05 MMTCO2 Eq
- Imported New 1990 estimate: 53.12 MMTCO2 Eq

#### Refineries

Includes emissions from fuel use for H2 production, on-site fuel combustion and some fugitive emissions

Data Source: California Energy Balance (PIIRA)


### Refineries - Updates

- Included:
  - H2 production emissions
  - Fugitive CH4 emissions
  - CH4 and N2O from fuel combustion
- Used IPCC oxidation factor of 1
- Extrapolated missing data for natural gas and associated liquids

#### Refineries - Estimates

CEC 1990 estimate:27.85 MMTCO2 Eq

New 1990 estimate: 30.45 MMTCO2 Eq



#### Landfills

- Includes emissions from decomposition of landfill material (CH4) and burning of landfill gas (CH4 and N2O)
- CO2 emissions from decomposition addressed in Land Use & Forestry
- Data Sources:
  - CIWMB for Waste in Place data and landfill controls
  - Survey data
  - IPCC, U.S.EPA and CIWMB for waste characterization
  - U.S.EPA for waste decay rate

#### Landfills - Updates

- Used IPCC Mathematically Exact First-Order Decay Model
- Tracks annual amount of carbon sequestered
  - Sequestered carbon not considered a sink
- Added N2O emissions

#### Landfills - Estimates

CEC 1990 estimate:8.13 MMTCO2 Eq



New 1990 estimate:6.58 MMTCO2 Eq

Landfill gas to energy estimate: 0.03 MMTCO2 Eq (included in electricity sector)

## Land Use, Land Use Change & Forestry (Biodegradable CO2)

- Includes CO2 emissions from:
  - Harvested and fuel wood
  - Landfilled and composted wood products
  - Biomass wildfires
  - Burning of crop residues (CO2 only)


- Includes removals from:
  - Forest, rangeland and crop growth

## Land Use, Land Use Change & Forestry - Updates

- Data Sources:
  - CEC Inventory (Winrock)
  - USFS and CIWMB data on wood consumption and wood landfill and composting data

- Used IPCC Atmospheric Flow Approach to track CO2 flux
- Removed landfill wood products as "sinks"

# Land Use, Land Use Change & Forestry - Estimates



CEC 1990 estimates (net):-17.16 MMTCO2 Eq

New 1990 estimates (net):-5.62 MMTCO2 Eq

## Transportation

- Includes CO2, CH4 and N2O emissions from:
  - Highway vehicles
  - Airplanes (domestic only)
  - Trains
  - Ships and Boats (domestic only)





## Highway Vehicles - Updates


- Data Sources:
  - California Energy Balance
  - EMFAC 2007

- Updates:
  - Used newer version of EMFAC for CH4
  - Updated diesel N2O emissions

## Highway Vehicles - Estimates

CEC 1990 estimate: 147.36 MMTCO2 Eq

New 1990 estimate: 139.81 MMTCO2 Eq



## Trains, Ships and Planes - Updates

- Data Sources:
  - California Energy Balance
  - U.S.EPA emission factors for CH4 and N2O

Updated CH4 and N2O emissions using U.S.EPA emission factors

## Trains, Ships and Planes - Estimates



CEC 1990 estimate:30.64 MMTCO2 Eq

New 1990 estimate: 30.76 MMTCO2 Eq

## Pending Updates & Improvements

- Cement fuel use
- Selected industrial processes (i.e., glass container manufacturing)
- Feedstock for hydrogen production for refineries
- Burning of associated gas and crude oil for oil and gas extraction
- Emissions from managed soils




## Documentation and Database

### Rationale for Documentation Efforts

- Requests by stakeholders
- Need for transparency

## Description of Database

- Created relational database
  - Stores equations, parameters and references
  - Calculates emissions
- Online access
  - Output available as PDF or Excel spreadsheet
  - Documentation pages



### Welcome to California

#### Cal/EPA Home ARB Home

#### Air Resources Board @

**ARB Links** 

DRAFT Documentation of California's Greenhouse Gas Inventory

Index last updated on August 07, 2007

Print Friendly

About ARB

**Programs** 

What's New

**Board Meetings** 

Calendar

**News Releases** 

Air Quality, Emissions & Modeling

Business Assistance & Training

Rulemakings

Laws & Regulations

Enforcement

**ARB Databases** 

Forms • Maps

Permits & Certifications, Etc.

Data documentation is a critical part of any inventory. This webpage provides links to draft documentation pages for the data and methods used in developing the statewide GHG inventory. Each link allows users to access information on estimated emissions, computations, activity data and other parameters for each year from 1990 to 2004. The categories in this index are based on those defined in 2006 by the Intergovernmental Panel on Climate Change (IPCC).

#### ■ — Index to the Documentation Pages —

- **± 1 ENERGY**
- 2 INDUSTRIAL PROCESSES AND PRODUCT USE
- ☐ 3 AGRICULTURE, FORESTRY, AND OTHER LAND USE
  - ☐ 3A Livestock
    - - 3A1a Cattle
      - ☐ 3A1c Sheep
        - Livestock population Sheep > CH4
      - ∃ 3A1d Goats
      - 3A1f Horses
        ■
      - 3A1h Swine
        3A1h Swine
  - 3B Land
  - ⊞ 3C Aggregate Sources and Non-CO2 Emissions Sources on Land
- # 4 WASTE



#### Air Resources Board



#### Documentation of California's Greenhouse Gas Inventory

[Back to Index] [Year 1990] [Previous year] [Next year] [Last year]

► Category: Enteric fermentation - Sheep

IPCC: 3A1c - Sheep

Sector: Agriculture & Forestry : Enteric Fermentation : Other Livestock : None

► Greenhouse gas: CH4

◆ 1990 CH4 from Livestock population - Sheep

Greenhouse gas estimation —

Amount: 8,000 metric tonnes of CH4 emitted (168,000 metric tonnes CO2eq.)

Basis: Calculation

Calculation: [Livestock population] \* [CH4 emission factor]

Reference: IPCC (2006) 2006 IPCC Guidelines for National Greenhouse Gas Inventories, Prepared by the National Greenhouse Gas Inventories Programmed, Eggleston HAS., Biennia L., Miwa K.,

Negara T. and Tanabe K. (eds). Vol.4, Chap. 10. Published: IGES, Japan.

Activity level —

Amount: Livestock population = 1,000,000 head of Sheep

Basis: Data

Reference: USDA (2006) U.S. Department of Agriculture, National Agriculture Statistics Service, California Statistics. Available online at: <a href="http://www.nass.usda.gov/Statistics.by">http://www.nass.usda.gov/Statistics.by</a> State/California/index.asp>

Parameters and Constants used in calculations —

Parameter: CH4 emission factor = 8000 g / individual

Reference: IPCC (2006) 2006 IPCC Guidelines for National Greenhouse Gas Inventories, Prepared by the National Greenhouse Gas Inventories Programmed, Eggleston HAS., Biennia L., Miwa K., Negara T. and Tanabe K. (eds). Vol.4, Chap. 10. Published: IGES, Japan.

Constant: Global warming potential of CH4 = 21

Reference: IPCC (1996). Second Assessment Report. Climate Change 1995: WG I - The Science of Climate Change. Intergovernmental Panel on Climate Change; J.T. Houghton, L.G. Meira Filho, B.A. Callander, N. Harris, A. Kattenberg, and K. Maskell (eds.); Cambridge University Press. Cambridge, U.K.

Greenhouse gas emitted per unit activity —

American C OO longer City and hand of Change



## 1990 Emissions Level

## Role of Inventory

- Statewide 1990 emissions level is basis for statewide, aggregated 2020 limit
- Starting point for Scoping Plan development

## Statewide 1990 Emissions Level

- Comprehensive of all anthropogenic emissions
- Excludes international bunker fuels (marine and aviation)

Aggregated number: 436 MMTCO2 Eq

## Statewide 1990 Emissions Level (cont.)

- Consider excluding domestic aviation
  - Equals approximately 24 MMTCO2 Eq

#### Rationale:

- State has no direct control over aviation-related fuel and emissions control technologies
- Interstate aviation akin to international aviation at statewide level

### 2020 Emissions Limit

- Equivalent to 1990 emissions level
- Statewide, aggregate emissions
- Limit not sector- or facility-specific
- Remains in effect unless otherwise amended or repealed

## 2020 Projections

- Ongoing work
- CAT projections based on 2003 IEPR
- CEC projections based on 2005 IEPR
- Draft 2007 IEPR under review
- Considering similar projection methodology to CEC

### Schedule

- Early Fall 2007
  - Draft staff report



- December 2007
  - Board consideration of GHG inventory and 1990 level / 2020 limit in public hearing

# Comments or Questions?



### **ARB Contacts**

Richard Bode – Chief Emissions Inventory Branch rbode@arb.ca.gov (916) 323-8413

Webster Tasat – Manager Emission Inventory Analysis Section wtasat@arb.ca.gov (916) 323-4950

Robert Jenne – Legal Counsel rjenne@arb.ca.gov (916) 322-3762

GHG Inventory Website <a href="http://www.arb.ca.gov/cc/ccei/ccei.htm">http://www.arb.ca.gov/cc/ccei/ccei.htm</a>



### **Staff Contacts**

Kevin Eslinger - Transportation

keslinge@arb.ca.gov (916) 445-2151

Larry Hunsaker – Electricity, Landfills, Refineries *Ihunsake@arb.ca.gov* (916) 324-7168

Neva Lowery - Agriculture nlowery@arb.ca.gov (916) 323-4887

Johnnie Raymond - Cement jraymond@arb.ca.gov (916) 324-7901

Jamesine Rogers – Staff Report jrogers@arb.ca.gov (916) 323-2722

Klaus Scott – Land Use, Land Use Change & Forestry kscott@arb.ca.gov (916) 327-0301

